arXiv:1108.2099v1 [math.KT] 10 Aug 2011

THE SPACE OF STABILITY CONDITIONS FOR QUIVERS WITH TWO VERTICES

TAKAHISA SHIINA

Abstract. The purpose of this article is to study the space of stability conditions $\mathrm{Stab}(P_n)$ on the bounded derived category $\mathcal{D}^b(P_n)$ of finite dimensional representations of a quiver P_n with two vertices and n parallel arrows. There is a local homeomorphism $\mathcal{Z} : \mathrm{Stab}(P_n) \to \mathbb{C}^2$. We show that, when the number of arrows is one or two, the map is a covering map if we restrict it to the complement of a line arrangement. When the number of arrows is greater than two we need to remove uncountably many lines to obtain a covering map.

1. Introduction

Tom Bridgeland introduced the notion of stability conditions on a triangulated category $\mathscr{T}$ ([Bri07]). A stability condition is a pair $(Z, \mathcal{P})$ of a group homomorphism Z from the Grothendieck group $K(\mathscr{T})$ to $\mathbb{C}$ and a family of abelian subcategories $\mathcal{P} = \{\mathcal{P}(\phi)\}_{\phi \in \mathbb{R}}$ of $\mathscr{T}$. The space $\mathrm{Stab}(\mathscr{T})$ of all (locally-finite) stability conditions on $\mathscr{T}$ can be made into a complex manifold. There is a continuous map $\mathcal{Z} : \mathrm{Stab}(\mathscr{T}) \to \mathrm{Hom}_{\mathbb{Z}}(K(\mathscr{T}), \mathbb{C})$ which maps $(Z, \mathcal{P})$ to Z. It is known that for each connected component $\Sigma \subset \mathrm{Stab}(\mathscr{T})$, there is a subspace $V(\Sigma) \subset \mathrm{Hom}_{\mathbb{Z}}(K(\mathscr{T}), \mathbb{C})$ such that $\mathcal{Z} : \Sigma \to V(\Sigma)$ is a local homeomorphism ([Bri07, Theorem 1.2]).

The spaces of stability conditions on the derived category of coherent sheaves on algebraic varieties have been studied intensively. For example, Bridgeland studied the space of stability conditions associated to a Kleinian singularity of ADE-type. Let A be a preprojective algebra of the ADE-graph Γ and let $\mathcal{D} = \mathcal{D}^b(\mathrm{modA})$. In this case, $\mathrm{Hom}_Z(K(\mathcal{D}), \mathbb{C})$ can be identified with the Cartan subalgebra $\mathfrak{h}$ of the Lie algebra $\mathfrak{g}$ corresponding to Γ. The image of $\mathcal{Z} : \mathrm{Stab}(\mathcal{D}) \to \mathfrak{h}$ is $\mathfrak{h}^{\mathrm{reg}}$, the complement of the affine root hyperplanes in $\mathfrak{h}$. Bridgeland conjectured that $\mathrm{Stab}(\mathcal{D})$ is connected and simply connected, and so $\mathcal{Z}$ is a covering map ([Bri09]). A. Ishii, K Ueda, and H. Uehara proved this conjecture in the case of type A ([IUU10]). However, the other cases of type D and E still remain open.

It is an important problem to find hyperplanes $\{H_\lambda\}_{\lambda \in \Lambda}$ in $\mathrm{Hom}_{\mathbb{Z}}(K(\mathscr{T}), \mathbb{C})$ such that the restriction of $\mathcal{Z}$ to the complement $\mathrm{Hom}_{\mathbb{Z}}(K(\mathscr{T}), \mathbb{C}) - \cup_{\lambda \in \Lambda} H_\lambda$ is a covering map.

There is a relation between the derived categories of finite dimensional representations of a quiver and the derived categories of algebraic varieties. For instance, the Kleinian singularities are related to the ADE-graphs and $\mathbb{P}^1$ is related to the Kronecker quiver. Emanuele Macrí studied the space of stability conditions on a triangulated category generated by finitely many exceptional objects ([Mac07]). A typical example of such a triangulated category is the derived category of a finite quiver. In particular, he proved $\mathrm{Stab}(P_n)$ is connected and simply connected where

---

Date: August 10, 2011.

$\mathrm{Stab}(P_n)$ denotes the space of stability conditions of the bounded derived category of finite dimensional representation of the quiver P_n with two vertices and n parallel arrows.

In this article, we give a concrete description of $\mathrm{Stab}(P_n)$ as a complex manifold. The local homeomorphism $\mathcal{Z} : \mathrm{Stab}(P_n) \to \mathrm{Hom}_{\mathbb{Z}}(K(P_n), \mathbb{C})$ is not a covering map. However, if we cut off some subspaces from $\mathrm{Hom}_{\mathbb{Z}}(K(P_n), \mathbb{C})$, the restricted map $\mathcal{Z}$ is a covering map. To see this the author use the $\mathbb{C}$-action on $\mathrm{Stab}(P_n)$ introduced by Okada and use the induced principal $\mathbb{C}$-bundle $\mathbb{C} \to \mathrm{Stab}(P_n) \to \mathrm{Stab}(P_n)/\mathbb{C}$.

Theorem 1.1. *The quotient space* $\mathrm{Stab}(P_n)/\mathbb{C}$ *is a* 1*-dimensional complex manifold given by gluing three, if* $n = 1$*, or countable, if* $n \geq 2$*, upper half planes* $\mathbb{H}_k$*. See Figure 1 and 2, where, in Figure 2, the sequence* $\{a_k\}$ *is the one defined in Definition 5.1.*

There is a map $\chi_n : \mathrm{Stab}(P_n)/\mathbb{C} \to \mathbb{C}\mathrm{P}^1$ *whose restriction onto* $\mathbb{C}\mathrm{P}^1 \setminus \{3\text{-points}\}$ $(n = 1)$*,* $\mathbb{C}\mathrm{P}^1 \setminus \{\text{countable points}\}$ $(n = 2)$*, or* $\mathbb{C}\mathrm{P}^1 \setminus \{\text{points and arc}\}$ $(n > 2)$ *is a covering map (see Figure 3, 4, and 5, respectively).*



FIGURE 1. $\mathrm{Stab}(P_1)/\mathbb{C}$


$\mathbb{H}_0$
$\mathbb{H}_k$
ψ_k
$\log \frac{a_{k-1}}{a_k}$
$\log \frac{a_k}{a_{k+1}}$
$\log \frac{a_{k+1}}{a_k}$
$\log \frac{a_k}{a_{k-1}}$
$\mathbb{H}_0$
$\mathbb{H}_{-k}$
ψ_{-k}
$\log \frac{a_{k+1}}{a_k}$
$\log \frac{a_k}{a_{k-1}}$
$\log \frac{a_{k-1}}{a_k}$
$\log \frac{a_k}{a_{k+1}}$

FIGURE 2. $\mathrm{Stab}(P_n)/\mathbb{C}$



FIGURE 3. χ_1



FIGURE 4. χ_2



FIGURE 5. χ_n

Theorem 1.2. *The local homeomorphism map* $\mathcal{Z}:\mathrm{Stab}(P_n)\to\mathrm{Hom}_{\mathbb{Z}}(K(P_n),\mathbb{C})$ *is a covering map if it is restricted to the complement of three lines* ($n=1$)*, countably many lines* ($n=2$)*, or a bundle of uncountably many lines* ($n>3$)*.*

The article is organized as follows. In section 2, we recall the notion of stability conditions introduced by Bridgeland, the action of universal cover of $\mathrm{GL}^+(2,\mathbb{R})$ on the space of stability conditions, and Macrí's results on the space of stability conditions on triangulated categories generated by finitely many exceptional objects. In section 3, we review Macrí's work on $\mathrm{Stab}(P_n)$. In section 4, we study $\mathrm{Stab}(P_1)$, i.e. the case of one arrow, and in section 5, $\mathrm{Stab}(P_n)$, the case of multiple arrows. In particular we examine the principal fiber bundle structure by $\mathbb{C}$-action, and examine how the coordinate neighborhoods are glued each other. In section 6 we prove main theorems.

Acknowledgements. I would like to thank Dai Tamaki for introducing me to the subject and invaluable advice.

2. STABILITY CONDITIONS

In this section we assume that a triangulated category $\mathscr{T}$ satisfies the following conditions.

- $\mathrm{Hom}^{\bullet}_{\mathscr{T}}(A,B) = \bigoplus_k \mathrm{Hom}^k_{\mathscr{T}}(A,B) = \bigoplus_k \mathrm{Hom}_{\mathscr{T}}(A,B[k])$ is a finite dimensional $\mathbb{C}$-vector space where $[k]$ is the k-fold shift functor.
- $\mathscr{T}$ is essentially small, that is the set of isomorphism classes of objects in $\mathscr{T}$ is finite,

Denote the Grothendieck group of $\mathscr{T}$ by $K(\mathscr{T})$ which is the quotient group of the free abelian group generated by all isomorphism classes of objects in $\mathscr{T}$ over the subgroup generated by the elements of the form $[A]+[B]-[C]$ for each distinguished triangle $A \to C \to B$.

2.1. **Definition of stability conditions.**

A *stability condition* on $\mathscr{T}$ is a pair $\sigma = (Z, \mathcal{P})$ of

- a group homomorphism $Z : K(\mathscr{T}) \to \mathbb{C}$ and
- full additive subcategories $\mathcal{P}(\phi)$ of $\mathscr{T}$ for each $\phi \in \mathbb{R}$

satisfying

Br1: if $\phi_1 > \phi_2$ and $A_j \in \mathcal{P}(\phi_j)$ then $\mathrm{Hom}_{\mathscr{T}}(A_1, A_2) = 0$,
Br2: $\mathcal{P}(\phi+1) = \mathcal{P}(\phi)[1]$,
Br3: each nonzero object has a Harder-Narasimhan filtration, and
Br4: if $0 \neq A \in \mathcal{P}(\phi)$, $Z([A]) = m(A)\exp(i\pi\phi)$ for some $m(A) \in \mathbb{R}_{>0}$.

A Harder-Narasimhan filtration of a nonzero object $E \in \mathscr{T}$ is a diagram

$$\begin{tikzcd}
0 = E_0 \arrow[rr] & & E_1 \arrow[r] \arrow[dl] & \cdots \arrow[r] & E_{n-1} \arrow[rr] & & E_n = E \arrow[dl] \\
& A_1 \arrow[ul, dotted] & & & & A_n \arrow[ul, dotted] &
\end{tikzcd}$$

such that each $E_{j-1} \to E_j \to A_j$ is a distinguished triangle in $\mathscr{T}$, $A_j \in \mathcal{P}(\phi_j)$ for all j and $\phi_1 > \cdots > \phi_n$. Note that for a stability condition $\sigma = (Z, \mathcal{P})$;

- a Harder-Narasimhan filtration for a nonzero object is unique up to isomorphism,
- Z is called the central charge,
- nonzero objects in $\mathcal{P}(\phi)$ is called semistable of phase ϕ,
- simple objects in $\mathcal{P}(\phi)$ is called stable,
- $\phi^+_\sigma(E) := \phi_1$, $\phi^-_\sigma(E) := \phi_n$, and
- $m_\sigma(E) := \sum |Z([A_j])|$ is called the mass of E.

A stability condition $(Z, \mathcal{P})$ is called locally-finite if there exists $\varepsilon \in \mathbb{R}$ such that $\mathcal{P}(\phi-\varepsilon, \phi+\varepsilon)$ is of finite length for each ϕ, that is, it is an Artinian and Noetherian category. Denote $\mathrm{Stab}(\mathscr{T})$ to be the set of all locally-finite stability conditions on $\mathscr{T}$. Bridgeland defined a generalized metric on $\mathrm{Stab}(\mathscr{T})$ by

$$d(\sigma_1, \sigma_2) = \sup_{0 \neq E \in \mathscr{T}} \left\{ |\phi^-_{\sigma_2}(E) - \phi^-_{\sigma_1}(E)|, |\phi^+_{\sigma_2}(E) - \phi^+_{\sigma_1}(E)|, \left| \log \frac{m_{\sigma_2}(E)}{m_{\sigma_1}(E)} \right| \right\} \in [0, \infty]$$

for $\sigma_1, \sigma_2 \in \mathrm{Stab}(\mathscr{T})$. It is proved that $\mathrm{Stab}(\mathscr{T})$ is locally homeomorphic to a complex vector space, so it is a complex manifold.

Theorem 2.1. [Bri07, Theorem 1.2] *For each connected component $\Sigma \subset \mathrm{Stab}(\mathscr{T})$ there are a linear subspace $V(\Sigma) \subset \mathrm{Hom}_{\mathbb{Z}}(K(\mathscr{T}), \mathbb{C})$, with a well-defined linear topology, and a local homeomorphism $\mathcal{Z} : \Sigma \to V(\Sigma)$ which maps a stability condition $(Z, \mathcal{P})$ to its central charge Z.*

Bridgeland proved that for any stability condition $(Z, \mathcal{P})$, the additive subcategory $\mathcal{P}(0, 1]$ generated by all semistables with phase in $(0, 1]$ is a heart of a bounded t-structure. Conversely, if a heart of a bounded t-structure and a stability function are given, a stability condition can be constructed. We denote $K(\mathcal{A})$ to be the Grothendieck group of an abelian category $\mathcal{A}$.

Definition 2.2. Let $\mathcal{A} \subset \mathscr{T}$ be a full abelian subcategory.

(1) A *stability function* on $\mathcal{A}$ is a group homomorphism $Z : K(\mathcal{A}) \to \mathbb{C}$ satisfying $Z([E]) \in H = \{m \exp(\pi i \phi) \,|\, m > 0,\ 0 < \phi \leq 1\}$ for all nonzero object $E \in \mathcal{A}$.

(2) For a stability function Z on $\mathcal{A}$, define $\phi(E)$ to be the phase of $Z([E])$.

(3) An object $E \in \mathcal{A}$ is said to be *semistable* if $\phi(F) \leq \phi(E)$ for each subobject $F \subset E$.

(4) A stability function Z on $\mathcal{A}$ is said to have the *Harder-Narasimhan property* if for every nonzero object $E \in \mathcal{A}$, there is a finite chain of subobjects
$$0 = E_0 \subset E_1 \subset \cdots \subset E_{n-1} \subset E_n = E$$
whose factors $F_j = E_j / E_{j-1}$ are semistable objects with
$$\phi(F_1) > \phi(F_2) > \cdots > \phi(F_n).$$

Theorem 2.3. [Bri07, Proposition 5.3] *To give a stability condition on a triangulated category $\mathscr{T}$ is equivalent to giving a bounded t-structure on $\mathscr{T}$ and a stability function on its heart with the Harder-Narasimhan property.*

2.2. **Group action on the space of stability conditions.** Bridgeland defined a right action of the group $\widetilde{\mathrm{GL}^+}(2, \mathbb{R})$, the universal cover of $\mathrm{GL}^+(2, \mathbb{R})$, on $\mathrm{Stab}(\mathscr{T})$ and a left action of $\mathrm{Aut}(\mathscr{T})$, the group of exact autoequivalences of $\mathscr{T}$, on $\mathrm{Stab}(\mathscr{T})$. Here these two actions commute.

Note that $\widetilde{\mathrm{GL}^+}(2, \mathbb{R})$ can be thought of as the set of pairs (T, f), where $f : \mathbb{R} \to \mathbb{R}$ is an increasing map with $f(\phi + 1) = f(\phi) + 1$ and $T \in \mathrm{GL}^+(2, \mathbb{R})$, such that $T(\exp(2\pi i \phi))/|T(\exp(2\pi i \phi))| = \exp(2\pi i f(\phi))$ in $\mathbb{C} = \mathbb{R}^2$ for all $\phi \in \mathbb{R}$. Then, for a stability condition $\sigma = (Z, \mathcal{P}) \in \mathrm{Stab}(\mathscr{T})$ and a pair $(T, f) \in \widetilde{\mathrm{GL}^+}(2, \mathbb{R})$, define a new stability condition $\sigma' = (Z', \mathcal{P}')$ by setting $Z' = T^{-1} \circ Z$ and $\mathcal{P}'(\phi) = \mathcal{P}(f(\phi))$.

Secondly note that an element $G \in \mathrm{Aut}(\mathscr{T})$ induces an automorphism g of $K(\mathscr{T})$. For $\sigma = (Z, \mathcal{P})$, define a new stability condition $G(\sigma)$ to be $(Z \circ g^{-1}, \mathcal{P}'')$ where $\mathcal{P}''(\phi) = G(\mathcal{P}(\phi))$.

Okada defined an action of $z = a + bi \in \mathbb{C}$ on $(Z, \mathcal{P}) \in \mathrm{Stab}(\mathscr{T})$ [Oka07, Definition 2.3] by $z \cdot (Z, \mathcal{P}) = (Z', \mathcal{P}')$ where $Z'(E) = e^z \cdot Z$ and $\mathcal{P}_z(\phi) = \mathcal{P}(\phi - \frac{b}{\pi})$.

Proposition 2.4. [Oka07, Proposition 4.1] *The $\mathbb{C}$-action is holomorphic and free. It coincides with the action of a subgroup of $\widetilde{\mathrm{GL}^+}(2, \mathbb{R})$. The quotient $\mathrm{Stab}(\mathscr{T})/\mathbb{C}$ is a complex manifold, modeled on the projective space of a topological vector space.*

2.3. **Stability conditions and exceptional objects.** Emanuele Macrì discussed the spaces of stability conditions on triangulated categories generated by finitely many exceptional objects [Mac07].

Definition 2.5. (1) An object E in $\mathscr{T}$ is called *exceptional* if

$$\mathrm{Hom}^k_{\mathscr{T}}(E,E) = \begin{cases} \mathbb{C} & (k=0) \\ 0 & (\text{otherwise}). \end{cases}$$

(2) A sequence $(E_1, E_2, \dots, E_n)$ is called an *exceptional collection* if each E_i is exceptional and $\mathrm{Hom}^k_{\mathscr{T}}(E_i, E_j) = 0$ for all k and $i > j$.
(3) An exceptional collection is called *complete* if it generates $\mathscr{T}$ by shifts and extensions.
(4) An exceptional collection is called *Ext-exceptional* if $\mathrm{Hom}^{\leq 0}(E_i, E_j) = 0$ for all $i \neq j$.

For an exceptional collection, a new exceptional collection can be constructed by mutations.

Definition 2.6. (1) Let E and F be exceptional objects. We define $\mathcal{L}_E F$ and $\mathcal{R}_F E$ by the following distinguished triangles:

$$\mathcal{L}_E F \longrightarrow \mathrm{Hom}^\bullet(E,F) \otimes E \longrightarrow F,$$

$$E \longrightarrow \mathrm{Hom}^\bullet(E,F)^* \otimes F \longrightarrow \mathcal{R}_F E.$$

We call $\mathcal{L}_E F$ *the left mutation* of F by E and $\mathcal{R}_F E$ *the right mutation* of E by F.

(2) Let $\mathcal{E} = (E_1, \dots, E_n)$ be an exceptional collection. Define $\mathcal{L}_i\mathcal{E}$ and $\mathcal{R}_i\mathcal{E}$ to be the sequences

$$\mathcal{L}_i\mathcal{E} = (E_1, \dots, E_{i-1}, \mathcal{L}_{E_i}E_{i+1}, E_i, E_{i+2}, \dots, E_n),$$

$$\mathcal{R}_i\mathcal{E} = (E_1, \dots, E_{i-1}, E_{i+1}, \mathcal{R}_{E_{i+1}}E_i, E_{i+2}, \dots, E_n).$$

We call $\mathcal{L}_i\mathcal{E}$ (resp. $\mathcal{R}_i\mathcal{E}$) *the left (resp. right) mutation* of $\mathcal{E}$.

Proposition 2.7. *(See for example* [Bon89]*)*

(1) *A mutation of an exceptional collection is also an exceptional collection.*
(2) *If an exceptional collection generates $\mathscr{T}$ then its mutation also generates $\mathscr{T}$.*
(3) *The following relations hold:*

$$\mathcal{L}_i\mathcal{R}_i = \mathcal{R}_i\mathcal{L}_i = id, \quad \mathcal{R}_i\mathcal{R}_{i+1}\mathcal{R}_i = \mathcal{R}_{i+1}\mathcal{R}_i\mathcal{R}_{i+1}, \quad \text{and} \quad \mathcal{L}_i\mathcal{L}_{i+1}\mathcal{L}_i = \mathcal{L}_{i+1}\mathcal{L}_i\mathcal{L}_{i+1}.$$

The last equations and, $\mathcal{R}_i\mathcal{R}_j = \mathcal{R}_j\mathcal{R}_i$ and $\mathcal{L}_i\mathcal{L}_j = \mathcal{L}_j\mathcal{L}_i$ if $|i-j| > 1$, induces an action of the braid group on the set of exceptional collections.

Here we denote by $\langle S \rangle$ the extension-closed subcategory of $\mathscr{T}$ generated by a subcategory S.

Lemma 2.8. [Mac07, Lemma 3.14] *Let $(E_1, \dots, E_n)$ be a complete Ext-exceptional collection on $\mathscr{T}$. Then $\langle E_1, \dots, E_n \rangle$ is the heart of a bounded t-structure on $\mathscr{T}$.*

Lemma 2.9. [Mac07, Lemma 3.16] *Let $(E_1, \dots, E_n)$ be a complete Ext-exceptional collection on $\mathscr{T}$ and let $(Z, \mathcal{P})$ be a stability condition on $\mathscr{T}$. Assume $E_1, \dots, E_n \in \mathcal{P}((0,1])$. Then $\langle E_1, \dots, E_n \rangle = \mathcal{P}((0,1])$ and E_j is stable, for all $j = 1, \dots, n$.*

Fix a complete exceptional collection $\mathcal{E} = (E_1, \dots, E_n)$ on $\mathscr{T}$. The Grothendieck group $K(\mathscr{T})$ is a free abelian group of finite rank isomorphic to $\mathbb{Z}^{\oplus n}$ generated by the classes $[E_i]$. By applying Theorem 2.3 and Lemma 2.8, we obtain the next Lemma.

Lemma 2.10. [Mac07] *A stability condition, as a pair $(Z, \mathcal{A})$ of a heart $\mathcal{A}$ and a stability function Z, can be constructed by the following steps:*

- *Choose integers $p_1, \dots, p_n$ such that $(E_1[p_1], \dots, E_n[p_n])$ is Ext-exceptional.*
- *Denote $\mathrm{Q}_p := \langle E_1[p_1], \dots, E_n[p_n] \rangle$. Then Q_p is the heart of a bounded t-structure (Lemma 2.8).*
- *Pick n points $z_1, \dots, z_n$ in the upper half plane H in $\mathbb{C}$ and define $Z_p : K(Q_p) \to \mathbb{C}$ by $Z_p(E_i[p_i]) = z_i$. Then Z is a stability function with the Harder-Narasimhan property.*

Define $\Theta_{\mathcal{E}}$ to be the subset of $\mathrm{Stab}(\underbrace{\mathscr{T})}$ consisting of stability conditions obtained by this way, up to the action of $\mathrm{GL}^+(2, \mathbb{R})$. Lemma 2.9 implies the following immediately.

Lemma 2.11. [Mac07] *For an exceptional collection $\mathcal{E} = (E_1, \dots, E_n)$, each E_i is stable for all stability conditions in $\Theta_{\mathcal{E}}$.*

Lemma 2.12. [Mac07, Lemma 3.19.] *The subspace $\Theta_{\mathcal{E}} \subset \mathrm{Stab}(\mathscr{T})$ is an open, connected and simply connected n-dimensional submanifold.*

This Lemma is one of the main results in [Mac07]. It is worth to note how Macrí proved this fact.

For an exceptional collection $\mathcal{F}_s = (F_1, \dots, F_s)$ $(s > 1)$, we define, for $i < j$,

$$k_{i,j}^{\mathcal{F}_s} := \begin{cases} +\infty, & \text{if } \mathrm{Hom}^k(F_i, F_j) = 0 \text{ for all k}, \\ \min\left\{ k \;\middle|\; \mathrm{Hom}^k(F_i, F_j) \neq 0 \right\}, & \text{otherwise}. \end{cases}$$

Then define $\alpha_s^{\mathcal{F}_s} = 0$, and for $i < s$,

$$\alpha_i^{\mathcal{F}_s} := \min_{j > i} \left\{ k_{i,j}^{\mathcal{F}_s} + \alpha_j^{\mathcal{F}_s} \right\} - (s - i - 1)$$

inductively. Consider $\mathbb{R}^n$ with coordinates $\phi_1, \dots, \phi_n$. Let $\mathcal{F}_s := (E_{\ell_1}, \dots, E_{\ell_s}) \subset (E_1, \dots, E_n)$, $s > 1$. Define the relation $R^{\mathcal{F}_s}$ on $\mathbb{R}^n$ by $\phi_{\ell_1} < \phi_{\ell_s} + \alpha_1^{\mathcal{F}_s}$. Finally define

$$\mathcal{C}_{\mathcal{E}} := \left\{ (m_1, \dots, m_n, \phi_1, \dots, \phi_n) \in \mathbb{R}^{2n} \;\middle|\; \begin{array}{l} m_i > 0 \text{ for all } i \text{ and} \\ R^{\mathcal{F}_s} \text{ for all } \mathcal{F}_s \subset \mathcal{E},\ s > 1 \end{array} \right\}.$$

Then $\mathcal{C}_{\mathcal{E}}$ is homeomorphic to $\Theta_{\mathcal{E}}$ via the map $\rho : \Theta_{\mathcal{E}} \to \mathcal{C}_{\mathcal{E}}$ defined by $m_i(\rho(\sigma)) := |Z(E_i)|$ and $\phi_i(\rho(\sigma)) := \phi_\sigma(E_i)$. To see [Mac07] for the proof of the fact that ρ is a homeomorphism and that $\mathcal{C}_{\mathcal{E}}$ is connected and simply connected.

3. QUIVERS WITH TWO VERTICES

A *quiver* Q consists of the following data;

- a set Q_0 of vertices,
- a set Q_1 of arrows,
- a map $s : Q_1 \to Q_0$ called the source map, and
- a map $t : Q_1 \to Q_0$ called the target map.

We write $\alpha : a \to b$ if $s(\alpha) = a$ and $t(\alpha) = b$ for $\alpha \in Q_1$. A *representation* V on Q is a family $V = \{V_a, f_\alpha\}$ such that

- V_a is a finite dimensional $\mathbb{C}$-vector space for all $a \in Q_0$ and
- $f_\alpha : V_a \to V_b$ is a linear map for each arrow $\alpha : a \to b$.

A *morphism* $F : V \to V'$ between representations is the family $F = \{F_a : V_a \to V'_a\}$ such that it is compatible with f_α. Define $\mathrm{rep}(Q)$ to be the abelian category of representations on Q and $\mathcal{D}(Q) = \mathcal{D}^b(\mathrm{rep}(Q))$ to be the bounded derived category of $\mathrm{rep}(Q)$. Here we consider quivers with no loops, no cycles and no relations. The following are well-known facts:

Q1: Define E_i in $\mathrm{rep}(Q)$ by assigning $\mathbb{C}$ to the vertex i and 0 to other vertices. Then $\{E_i\}_{i \in Q_0}$ is a complete set of simple objects.

Q2: $\mathrm{Hom}(S_i, S_j) = \mathbb{C}$ if $i = j$ and $= 0$ if $i \neq j$.

Q3: $\mathrm{Ext}^1(S_i, S_j) = \mathbb{C}^n$ if there exist n arrows from i to j.

Q4: $\mathrm{rep}(Q)$ is hereditary. And so every $X \in \mathcal{D}(Q)$ is isomorphic to the its cohomology considered as a complex with 0 differential. (See for example [Kra07, 1.6].)

Q5: $K(\mathcal{D}(Q)) \cong K(\mathrm{rep}(Q))$ is a free abelian group generated by all classes $[E_i]$.

As a triangulated category, $\mathcal{D}(Q)$ fits into the settings of the previous section and Macrí's results can be applied to $\mathcal{D}(Q)$. In fact Macrí studied $\mathrm{Stab}(P_n) = \mathrm{Stab}(\mathcal{D}(P_n))$ where P_n is the quiver with two vertices and n parallel arrows from one vertex to the other.

$$\begin{tikzcd}P_n : \overset{0}{\bullet} \arrow[r, shift left=2] \arrow[r] \arrow[r, shift right=2] \arrow[r, phantom, "\vdots"] & \overset{1}{\bullet}\end{tikzcd}$$

Let S_0 and S_1 be the exceptional objects in $\mathcal{D}(P_n)$ where $S_0[1]$ and S_1 are the simple objects in $\mathrm{rep}(P_n)$. According to [Mac07, CB92, Rin94], if we define

$$S_k := \begin{cases} \mathcal{L}_{S_{k+1}} S_{k+2} & (k < 0) \\ \mathcal{R}_{S_{k-1}} S_{k-2} & (k \geq 2) \end{cases}$$

inductively, then S_k's are the only exceptional objects in $\mathcal{D}(P_n)$ up to shifts and isomorphisms and each adjacent pair (S_k, S_{k+1}) is an exceptional collection. Note that each (S_k, S_{k+1}) is the right mutation of (S_{k-1}, S_k). Note also that if $n = 1$, iterating mutation tree times is equal to the degree one shift;

$$\mathcal{R}_1 \mathcal{R}_1 \mathcal{R}_1 (S_0, S_1) = (S_0[1], S_1[1])$$

and so S_0, S_1 and S_2 are the only essentially different exceptional objects.

Lemma 3.1. [Mac07, Lemma 4.1] *Assume that* $n > 1$, *if* $i < j$ *then in* $\mathcal{D}(P_n)$

- $\mathrm{Hom}^k(S_i, S_j) \neq 0$ *only if* $k = 0$;
- $\mathrm{Hom}^k(S_j, S_i) \neq 0$ *only if* $k = 1$.

In particular the pair (S_k, S_{k+1}) *is a complete strong exceptional collection.*

Lemma 3.2. [Mac07, Lemma 4.2] *In every stability condition on* $\mathcal{D}(P_n)$ *there exists a stable exceptional pair* (E, F).

Let Θ_k, $k \in \mathbb{Z}$, be a subset of $\mathrm{Stab}(P_n)$ consisting of all stability conditions made from (S_k, S_{k+1}) by Lemma 2.10.

Lemma 3.3. Θ_k *coincides with the subset of* $\mathrm{Stab}(P_n)$ *consisting of all stability conditions in which* S_k *and* S_{k+1} *are stable.*

Proof. See [Mac07, section 4]. $\square$

Proposition 3.4. *If* $n > 1$, $\mathrm{Stab}(P_n) = \bigcup_{k\in\mathbb{Z}} \Theta_k$. *In addition,* $\mathrm{Stab}(P_1) = \Theta_0 \cup \Theta_1 \cup \Theta_2$

Proof. It follows immediately from Lemma 3.1, 3.2 and 3.3. $\square$

Proposition 3.5. [Mac07, Proposition 4.4] *For all integers* $k \neq h$ *we have*

$$\Theta_k \cap \Theta_h = O_{-1}$$

where O_{-1} *is the* $\widetilde{\mathrm{GL}^+(2,\mathbb{R})}$*-orbit of the stability condition* $\sigma_{-1} = (Z_{-1}, \mathcal{P}_{-1})$ *given by* $Z_{-1}(S_0[1]) = -1$ *and* $Z_{-1}(S_1) = 1 + i$.

According to the proof of Lemma 2.12, we have

$$\Theta_k \cong \mathcal{C}_k := \{(m_1, m_2, \phi_1, \phi_2) \in \mathbb{R}^4 \mid m_i > 0 \text{ and } \phi_1 < \phi_2\}.$$

Clearly, it is connected and simply connected. By using the Seifert-Van Kampen theorem, Macrí shows that the whole space of stability conditions is connected and simply connected.

Theorem 3.6. [Mac07, Theorem 4.5] $\mathrm{Stab}(P_n)$ *is a connected and simply connected* 2*-dimensional complex manifold.*

In the rest of this article, we analyze how Θ_k's are glued each other. For convenience, let us change the expression of $\mathcal{C}_k$.

Lemma 3.7. $\mathcal{C}_k \cong \mathbb{C} \times \mathbb{H}$ *where* $\mathbb{H} = \{z \in \mathbb{C} \mid \mathcal{I}m\, z > 0\}$*, the upper half plane.*

Proof. Define

$$\zeta_k(m_1, m_2, \phi_1, \phi_2) := \left(\log(m_1) + i\pi\phi_1,\, \log\left(\frac{m_2}{m_1}\right) + i\pi(\phi_2 - \phi_1)\right).$$

It is clear that ζ_k is a homeomorphism. $\square$

Let us consider the correspondence between elements in Θ_k and $\mathbb{C} \times \mathbb{H}$. For $z = a + bi \in \mathbb{C}$ and $w = c + di \in \mathbb{H}$, $\sigma = (Z, \mathcal{P}) \in \Theta_k$ is the one defined by

$$Z(S_k) = e^z,\ Z(S_{k+1}) = e^{z+w},\ \phi_\sigma(S_k) = b \ \text{ and } \ \phi_\sigma(S_{k+1}) = b + d. \tag{1}$$

Conversely, for $\sigma = (Z, \mathcal{P}) \in \Theta_k$,

$$\begin{aligned} z &= \log|Z(S_k)| + i\pi\phi_\sigma(S_k) \ \text{ and} \\ w &= \log\left|\frac{Z(S_{k+1})}{Z(S_k)}\right| + i\pi(\phi_\sigma(S_{k+1}) - \phi_\sigma(S_k)). \end{aligned}$$

Lemma 3.8. *The action of* $\mathbb{C}$ *on* $\mathrm{Stab}(P_n)$*, defined by Okada, can be restricted to* Θ_k *for each* k.

Proof. The $\mathbb{C}$-action does not change the stable objects. $\square$

The induced $\mathbb{C}$ action on $\mathbb{C} \times \mathbb{H}$ is given by

$$z' \cdot (z, w) = (z' + z, w). \tag{2}$$

Therefore the quotient is homeomorphic to $\mathbb{H}$.

4. THE CASE OF ONE ARROW

In this section, we study $\mathrm{Stab}(P_1)$. In this case, since essentially different exceptional collections in $\mathcal{D}(P_1)$ are only three: (S_0, S_1), (S_1, S_2) and $(S_2, S_3 = S_0[1])$, the space of stability conditions is union of subsets constructed by them:

$$\mathrm{Stab}(P_1) = \Theta_0 \cup \Theta_1 \cup \Theta_2.$$

The intersection of Θ_i's is $O_{-1} \subset \mathrm{Stab}(P_1)$ consisting of all stability conditions such that S_0, S_1 and S_2 are all stable. Since there is a distinguished triangle

$$S_0 \longrightarrow S_1 \longrightarrow S_2 \longrightarrow S_0[1],$$

we have

$$Z(S_1) = Z(S_0) + Z(S_2) \tag{3}$$

and

$$\phi_\sigma(S_0) < \phi_\sigma(S_1) < \phi_\sigma(S_2) < \phi_\sigma(S_0) + 1 \tag{4}$$

for all $\sigma = (Z, \mathcal{P}) \in O_{-1}$.



FIGURE 6.

By Lemma 3.7, let $\Theta_k \cong \mathbb{C}_k \times \mathbb{H}_k$ $(k = 0, 1, 2)$ where $\mathbb{C}_k$ and $\mathbb{H}_k$ are copies of $\mathbb{C}$ and $\mathbb{H}$. Now we consider how $\mathbb{C}_k \times \mathbb{H}_k$ are glued each other.

Theorem 4.1. *Let*

$$\overline{\mathbb{H}}_k := \{w \in \mathbb{H} \,|\, 0 < \mathcal{I}m\, w < 1\}.$$

Define $\varphi_k : \mathbb{C}_0 \times \overline{\mathbb{H}}_0 \to \mathbb{C}_k \times \overline{\mathbb{H}}_k$ *$(k = 1, 2)$ by*

$$\varphi_1(z, w) := \left(z + w,\, \log \frac{e^w - 1}{e^w} \right)$$

$$\varphi_2(z, w) := \left(z + \log(e^w - 1),\, \log \frac{1}{1 - e^w} \right)$$

and define an equivalence relation by $(z, w) \sim \varphi_k(z, w)$ *for each* $(z, w) \in \mathbb{C}_0 \times \overline{\mathbb{H}}_0$ *and* k*. Then*

$$\mathrm{Stab}(P_1) \cong \left(\bigcup_{k=0}^{2} \mathbb{C}_k \times \mathbb{H}_k \right) / \sim$$

Proof. Because of the inequality (4), Θ, the intersection of Θ_k's, corresponds to $\mathbb{C}_k \times \overline{\mathbb{H}}_k$ for each k. Let $(Z, \mathcal{P}) \in \Theta$ and let $(z_k, w_k) \in \mathbb{C}_k \times \overline{\mathbb{H}}_k$ ($k = 0,\ 1,\ 2$) be elements corresponding to $(Z, \mathcal{P})$. The equation (1), in section 3, and Figure 6 imply that

$$e^{z_1} = Z(S_1) = e^{z_0+w_0} \quad \text{and} \quad e^{z_1+w_1} = Z(S_2) = e^{z_0+w_0} - e^{z_0}$$

and

$$e^{z_2} = Z(S_2) = e^{z_0+w_0} - e^{z_0} \quad \text{and} \quad e^{z_2+w_2} = Z(S_0[1]) = -e^{z_0}.$$

Hence φ_k are defined.

It is clear that φ_i are homeomorphisms and Θ_i's are glued by them. ☐

Theorem 4.2. *Define* $\psi_k : \overline{\mathbb{H}}_0 \to \overline{\mathbb{H}}_k$ *(*$k = 1, 2$*) by*

$$\psi_1(w) := \log \frac{e^w - 1}{e^w} \quad \textit{and} \quad \psi_2(w) := \log \frac{1}{1 - e^w}$$

define an equivalence relation by $w \sim \psi_k(w)$ *for each* $w \in \overline{\mathbb{H}}_0$ *and* k*, and define*

$$\mathcal{C}_1 := \left(\bigcup_{k=0}^{2} \mathbb{H}_k \right) / \sim\ .$$

Then $\mathrm{Stab}(P_1)/\mathbb{C} \cong \mathcal{C}_1$ *(see Figure 1).*

Proof. The action of $\mathbb{C}$ on $\mathbb{C}_k \times \mathbb{H}_k$ is given by the addition to the first factor, c.f. (2) in section 3. Hence the quotients $\mathbb{H}_k$ by $\mathbb{C}$ are glued by ψ_k, which is the second factor of φ_k. ☐

5. THE CASE OF MULTIPLE ARROWS

Suppose $n \geq 2$. On the bounded derived category $\mathcal{D}(P_n)$ of representations of P_n, there is a sequence of exceptional objects

$$\dots, S_{-2},\ S_{-1},\ S_0,\ S_1,\ S_2,\ \dots$$

where each pair (S_k, S_{k+1}) is a complete strong exceptional pair, each S_{k+2} is a right mutation of $S_k \to S_{k+1}$, and no other exceptional objects without shifts and isomorphisms.

Definition 5.1. For $n \geq 2$, define a sequence of real numbers $\{a_k\}_{k \in \mathbb{Z}}$ by

- $a_0 = 0$, $a_1 = 1$,
- $a_{k+2} = na_{k+1} - a_k$ for $k \geq 2$, and
- $a_{-\ell} = -a_\ell$ for $\ell \geq 1$.

Lemma 5.2. *There is a distinguished triangle*

$$\cdots \longrightarrow S_{k-2} \longrightarrow S_{k-1}^{\oplus n} \longrightarrow S_k \longrightarrow S_{k-2}[1] \longrightarrow \cdots$$

Hence, for a group homomorphism $Z : K(P_n) \to \mathbb{C}$*, we obtain*

$$Z(S_k) = a_k Z(S_1) - a_{k-1} Z(S_0). \tag{5}$$

Proof. It is well known that

$$\mathrm{Hom}^j_{\mathcal{D}^b(P_n)}(S_0, S_1) \cong \begin{cases} \mathbb{C}^n & (j = 0) \\ 0 & (\text{otherwise}). \end{cases}$$

By the definition of mutation, there is a triangle

$$\cdots \longrightarrow S_0 \longrightarrow \mathrm{Hom}^\bullet_{\mathcal{D}^b(P_n)}(S_0, S_1) \otimes S_1 = S_1^{\oplus n} \longrightarrow S_2 \longrightarrow S_0[1] \longrightarrow \cdots .$$

The Hom exact sequence implies

$$\operatorname{Hom}^j_{\mathcal{D}^b(P_n)}(S_1, S_2) \cong \begin{cases} \mathbb{C}^n & (j=0) \\ 0 & (\text{otherwise}) \end{cases}$$

and

$$\operatorname{Hom}^j_{\mathcal{D}^b(P_n)}(S_2, S_0) \cong \begin{cases} \mathbb{C} & (j=1) \\ 0 & (\text{otherwise}). \end{cases}$$

We can show a similar property holds for all S_k by induction.

For a group homomorphism $Z : K(P_n) \to \mathbb{C}$, the triangle induces

$$\begin{aligned} Z(S_k) &= nZ(S_{k-1}) - Z(S_{k-2}) \qquad (\text{for all } k>1 \text{ and}) \\ Z(S_{-k}) &= nZ(S_{-k+1}) - Z(S_{-k+2}) \qquad (\text{for all } k>0.) \end{aligned}$$

By induction we obtain the equation (5). □

The sequence $\{a_k\}$ can be described as follows. If $n=2$

$$a_0 = 0,\ a_1 = 1,\ a_2 = 2,\ \dots,\ a_k = k, \dots,$$

and if $n>2$

$$\begin{aligned} &a_0 = 0,\ a_1 = 1,\ a_2 = n,\ a_3 = n^2-1, \dots, \\ &a_k = \frac{1}{\sqrt{n^2-4}} \left\{ \left(\frac{n+\sqrt{n^2-4}}{2} \right)^k - \left(\frac{n-\sqrt{n^2-4}}{2} \right)^k \right\}, \dots . \end{aligned}$$

Note that, for $n \geq 2$, $\left\{\frac{a_{k+1}}{a_k}\right\}$ and $\left\{\frac{a_k}{a_{k+1}}\right\}$ are convergent sequences.

$$\begin{aligned} &\frac{a_2}{a_1} > \frac{a_3}{a_2} > \cdots > \frac{a_k}{a_{k-1}} > \frac{a_{k+1}}{a_k} > \cdots \xrightarrow{k\to\infty} \frac{n+\sqrt{n^2-4}}{2} \\ &\frac{a_1}{a_2} < \frac{a_2}{a_3} < \cdots < \frac{a_{k-1}}{a_k} < \frac{a_k}{a_{k+1}} < \cdots \xrightarrow{k\to\infty} \frac{n-\sqrt{n^2-4}}{2} \end{aligned} \tag{6}$$

The space of stability conditions can be decomposed as

$$\operatorname{Stab}(P_n) \cong \bigcup_{k\in\mathbb{Z}} \Theta_k\,,$$

where each Θ_k is the subset of all stability conditions with S_k and S_{k+1} stable and is homeomorphic to $\mathbb{C}_k \times \mathbb{H}_k$. The intersection $O_{-1} = \Theta_k \cap \Theta_h$ (for all $k \neq h$) is the subset of all stability conditions on which all S_k are stable. Let us investigate how Θ_k are glued by the method used in the previous section.

Theorem 5.3. *Define* $\varphi_k : \mathbb{C}_0 \times \overline{\overline{\mathbb{H}}}_0 \to \mathbb{C}_k \times \overline{\overline{\mathbb{H}}}_k$ *by*

$$\varphi_k(z,w) := \left(z + \log(a_k e^w - a_{k-1}),\ \log \frac{a_{k+1}e^w - a_k}{a_k e^w - a_{k-1}} \right) \tag{7}$$

for each k *and define an equivalence relation by* $(z,w) \sim \varphi_k(z,w)$ *for all* $(z,w) \in \mathbb{C}_0 \times \overline{\overline{\mathbb{H}}}_0$ *and* k*. Then*

$$\operatorname{Stab}(P_n) \cong \left(\bigcup_{k\in\mathbb{Z}} \mathbb{C}_k \times \mathbb{H}_k \right) / \sim\ .$$

Proof. Because of Lemma 5.2 (5), O_{-1} corresponds to $\mathbb{C}_k \times \overline{\mathbb{H}}_k$ for each k. Let $(Z, \mathcal{P}) \in O_{-1}$, and let $(z, w) \in \mathbb{C}_0 \times \overline{\mathbb{H}}_0$ and $(z', w') \in \mathbb{C}_k \times \overline{\mathbb{H}}_k$ be elements corresponding to $(Z, \mathcal{P})$. From (1), we have $Z(S_0) = e^z$, $Z(S_1) = e^{z+w}$, $Z(S_k) = e^{z'}$, and $Z(S_{k+1}) = e^{z'+w'}$. Hence, the equation and Lemma 5.2 (5) imply

$$e^{z'} = Z(S_k) = a_k e^{z+w} - a_{k-1} e^z \ , \ e^{z'+w'} = \frac{Z(S_{k+1})}{Z(S_k)} = \frac{a_{k+1} e^{z+w} - a_k e^z}{a_k e^{z+w} - a_{k-1} e^z} \, .$$

By solving the equations, we obtain φ_k. □

Theorem 5.4. *Define* $\psi_k : \overline{\mathbb{H}}_0 \to \overline{\mathbb{H}}_k$ *by*

$$\psi_k(w) := \log \frac{a_{k+1} e^w - a_k}{a_k e^w - a_{k-1}}$$

for each k*, define an equivalence relation by* $w \sim \psi_k(w)$ *for each* $w \in \overline{\mathbb{H}}_0$ *and* k*, and define*

$$\mathcal{C}_n := \left(\bigcup_{k \in \mathbb{Z}} \mathbb{H}_k \right) / \sim \ .$$

Then $\mathrm{Stab}(P_n)/\mathbb{C} \cong \mathcal{C}_n$ *(see Figure 2).*

Proof. Similar to the proof of Theorem 4.2. □

6. PROOF OF MAIN THEOREMS

We have seen details of the principal $\mathbb{C}$-bundle structure of $\mathrm{Stab}(P_n)$. In this section we study the local homeomorphism $\mathcal{Z} : \mathrm{Stab}(P_n) \to \mathrm{Hom}_{\mathbb{Z}}(K(P_n), \mathbb{C})$.

Since the Grothendieck group of the derived category of representations of a quiver is a free abelian group generated by the isomorphism classes of simple objects, $K(P_n) \cong \mathbb{Z}\langle S_0, S_1 \rangle$. It is clear that $\mathrm{Hom}_{\mathbb{Z}}(K(P_n), \mathbb{C}) \cong \mathbb{C}^2$ by mapping Z to $(Z(S_0), Z(S_1))$.

Lemma 6.1. *The image of* $\mathcal{Z} : \mathrm{Stab}(P_n) \to \mathrm{Hom}_{\mathbb{Z}}(K(P_n), \mathbb{C})$ *is homeomorphic to* $\mathbb{C}^2 \setminus \{(0, 0)\}$.

Proof. It is enough to define a stability function Z, on some abelian subcategory $\langle S_k[p_k], S_{k+1}[p_{k+1}] \rangle$ generated by an Ext-exceptional pair, such that $Z([S_i]) = z_i$ $(i = 0, 1)$ for each $(z_0, z_1) \in \mathbb{C}^2 \setminus \{(0, 0)\}$.

Suppose z_0 and z_1 are both not equal to zero. We can choose $\varepsilon_i \in \{1, -1\}$ $(i = 0, 1)$ such that $\varepsilon_i z_i$ are both in the upper half plane. Then we can define a stability function Z on $\langle S_0[p_0], S_1[p_1] \rangle$ by $Z(S_i[p_i]) = \varepsilon_i z_i$ for a suitable choice of (p_0, p_1).

Suppose $z_0 = 0$ and $z_1 \neq 0$. We can choose $\varepsilon \in \{1, -1\}$ such that εz_1 is in the upper half plane. Then we can define a stability function Z on $\langle S_1[p_1], S_2[p_2] \rangle$ by $Z(S_i[p_i]) = \varepsilon z_1$ for a suitable choice of (p_1, p_2).

If $z_1 = 0$ and $z_0 \neq 0$, we can define Z on $\langle S_{-1}[p_{-1}], S_0[p_0] \rangle$ similarly. □

For the subset Θ_k consisting of all stability conditions with stable S_k and S_{k+1}, there is a commutative diagram

$$\begin{tikzcd}
\Theta_k \arrow[r, "f", "\cong"'] \arrow[d, "\mathcal{Z}"'] & \mathbb{C}_k \times \mathbb{H}_k \arrow[d, "g_k"] \\
\mathrm{Hom}_{\mathbb{Z}}(K(P_n), \mathbb{C}) \arrow[r, "h", "\cong"'] & \mathbb{C}^2
\end{tikzcd}$$

where $f(Z,\mathcal{P})=(z,w)$ defined by the equation (1). Since h is a homeomorphism given by $h(Z)=(Z(S_0),Z(S_1))$, the equality (5) implies

$$h(Z)=\left(\frac{a_kZ(S_{k+1})-a_{k+1}Z(S_k)}{a_{k-1}a_{k+1}-a_k^2},\frac{a_{k-1}Z(S_{k+1})-a_kZ(S_k)}{a_{k-1}a_{k+1}-a_k^2}\right)$$

and

$$g_k(z,w)=\left(\frac{a_ke^{z+w}-a_{k+1}e^z}{a_{k-1}a_{k+1}-a_k^2},\frac{a_{k-1}e^{z+w}-a_ke^z}{a_{k-1}a_{k+1}-a_k^2}\right).$$

It is clear that g_k commutes with the $\mathbb{C}$-action on $\mathbb{C}_k\times\mathbb{H}_k$ and the $\mathbb{C}^\times=\mathbb{C}\setminus\{0\}$-action on $\mathbb{C}^2$, because the $\mathbb{C}$-action is given by the addition to the first factor. It is also clear that $g_k\circ\varphi_k(z,w)=(e^z,e^{z+w})$. Therefore we obtain the following commutative diagram.

$$\begin{tikzcd}
\mathbb{C} \arrow[r] \arrow[d, "\exp"'] & \bigcup_{k\in\mathbb{Z}}\mathbb{C}_k\times\mathbb{H}_k/\sim \arrow[r] \arrow[d, "g"'] & \bigcup_{k\in\mathbb{Z}}\mathbb{H}_k/\sim \arrow[d, "\chi_n"'] \\
\mathbb{C}^\times \arrow[r] & \mathbb{C}^2\setminus\{(0,0)\} \arrow[r, "\pi"] & \mathbb{CP}^1
\end{tikzcd}$$

This shows the theorem:

Theorem 6.2. *We obtain the following commutative diagram where each rows are principal fiber bundles:*

$$\begin{tikzcd}
\mathbb{C} \arrow[r] \arrow[d, "\exp"'] & \mathrm{Stab}(P_n) \arrow[r] \arrow[d, "\mathcal{Z}"'] & \mathcal{C}_n \arrow[d, "\chi_n"'] \\
\mathbb{C}^\times \arrow[r] & \mathbb{C}^2\setminus\{(0,0)\} \arrow[r, "\pi"] & \mathbb{CP}^1
\end{tikzcd}$$

Now we consider the map χ_n. For $(z,w)\in\mathbb{C}_k\times\mathbb{H}_k$, the composition of g_k and π induces

$$\pi\circ g_k(z,w)=\left[\frac{a_ke^{z+w}-a_{k+1}e^z}{a_{k-1}a_{k+1}-a_k^2}:\frac{a_{k-1}e^{z+w}-a_ke^z}{a_{k-1}a_{k+1}-a_k^2}\right].$$

So we obtain

$$\chi_n(w)=[a_ke^w-a_{k+1}:a_{k-1}e^w-a_k]$$

for $w\in\mathbb{H}_k$. Note that $\chi_n(w)=[1:e^w]$ if $w\in\mathbb{H}_0$.

Proof of Theorem 1.1. In the case of $n=1$, since

$$\begin{aligned}h(Z)&=(Z(S_0),Z(S_1))\\&=(Z(S_1)-Z(S_2),Z(S_1))\\&=(-Z(S_0[1]),Z(S_2)-Z(S_0[1]))\end{aligned}$$

(see Figure 6), it can be easily seen that

$$\begin{aligned}g_1(z_1,w_1)&=(e^{z_1}-e^{z_1+w_1},e^{z_1})\\g_2(z_2,w_2)&=(-e^{z_2+w_2},e^{z_2}-e^{z_2+w_2})\end{aligned}$$

where $g_k:\mathbb{C}_k\times\mathbb{H}_k\to\mathbb{C}^2$. Hence

$$\chi_1(w)=\begin{cases}[1:e^w] & (w\in\mathbb{H}_0)\\ [1-e^w:1] & (w\in\mathbb{H}_1)\\ [e^w:e^w-1] & (w\in\mathbb{H}_2).\end{cases}$$

The projective special linear group $\mathrm{PSL}(2,\mathbb{C})$ acts on the sphere $\mathbb{C}\mathrm{P}^1$. Let $G_1 = \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix} \in \mathrm{PSL}(2,\mathbb{C})$, the order of which is 3. It is clear that

$$G_1 \cdot [1 : e^w] = [e^w : e^w - 1] \quad \text{and} \quad G_1^2 \cdot [1 : e^w] = [1 - e^w : 1],$$

that is $\chi_1(\mathbb{H}_1) = G_1^2 \cdot \chi_1(\mathbb{H}_0)$ and $\chi_1(\mathbb{H}_2) = G_1 \cdot \chi_1(\mathbb{H}_0)$. It is also clear that

$$\begin{cases} G_1 \cdot [1:0] = [0:1] \\ G_1 \cdot [0:1] = [1:1] \end{cases} \quad \text{and} \quad \begin{cases} G_1^2 \cdot [1:0] = [1:1] \\ G_1^2 \cdot [0:1] = [1:0] \end{cases}.$$

Therefore χ_1 wraps each upper half plane $\mathbb{H}_k$ on $\mathbb{C}\mathrm{P}^1$, i.e.

- wraps $\mathbb{H}_0$ on $\mathbb{C}\mathrm{P}^1$ around points $[0 : 1]$ and $[1 : 0]$,
- wraps $\mathbb{H}_1$ on $\mathbb{C}\mathrm{P}^1$ around points $[1 : 1]$ and $[1 : 0]$, and
- wraps $\mathbb{H}_2$ on $\mathbb{C}\mathrm{P}^1$ around points $[1 : 1]$ and $[0 : 1]$.

See Figure 3 for detail. Moreover we have thus proved that χ_1 is a covering map if it is restricted to the inverse image of

$$\mathcal{CP}_1 = \mathbb{C}\mathrm{P}^1 \setminus \{[1:0],\, [0:1],\, [1:1]\}. \tag{8}$$

In the case of $n = 2$, since $a_k = k$,

$$\chi_2(w) = [ke^w - (k+1) : (k-1)e^w - k] \quad \text{if } w \in \mathbb{H}_k.$$

Let $G_2 = \begin{bmatrix} 0 & 1 \\ -1 & 2 \end{bmatrix} \in \mathrm{PSL}(2,\mathbb{C})$, the power of which is

$$G_2^k = \begin{bmatrix} -k+1 & k \\ -k & k+1 \end{bmatrix}$$

for each $k \in \mathbb{Z}$. It is clear that

$$\begin{aligned} G_2^{-k} \cdot [1 : e^w] &= [(k+1) - ke^w : k + (-k+1)e^w] \\ &= [ke^w - (k+1) : (k-1)e^w - k], \end{aligned}$$

that is $\chi_2(\mathbb{H}_k) = G_2^{-k} \cdot \chi_2(\mathbb{H}_0)$. It is also clear that

$$\begin{cases} G_2^{-k} \cdot [1:0] = [k+1 : k] \quad \text{and} \\ G_2^{-k} \cdot [0:1] = [-k : -k+1] = [k : k-1]. \end{cases}$$

Therefore χ_2 wraps

- $\mathbb{H}_0$ on $\mathbb{C}\mathrm{P}^1$ around the axis from $[0 : 1]$ to $[1 : 0]$ and
- $\mathbb{H}_k$ on $\mathbb{C}\mathrm{P}^1$ around the axis from $[k : k-1]$ to $[k+1 : k]$.

Note that

$$\begin{aligned} \lim_{k\to\infty} [k : k-1] &= \lim_{k\to\infty} [k+1 : k] = [1 : 1] \\ \lim_{k\to\infty} [k-1 : k] &= \lim_{k\to\infty} [k : k+1] = [1 : 1]. \end{aligned}$$

See Figure 4. We have thus proved that χ_2 is a covering map if it is restricted to

$$\mathcal{CP}_2 = \mathbb{C}\mathrm{P}^1 \setminus \{[1:1],\, [k : k+1] \mid k \in \mathbb{Z}\}. \tag{9}$$

In the case of $n > 3$, let $G_n = \begin{bmatrix} 0 & 1 \\ -1 & n \end{bmatrix} \in \mathrm{PSL}(2,\mathbb{C})$, the power of which is

$$G_n^k = \begin{bmatrix} -a_{k-1} & a_k \\ -a_k & a_{k+1} \end{bmatrix}$$

for all $k \in \mathbb{Z}$. It is clear that

$$\begin{aligned} G_n^{-k} \cdot [1 : e^w] &= [-a_{-k-1} + a_{-k} e^w : -a_{-k} + a_{-k+1} e^w] \\ &= [a_{k+1} - a_k e^w : a_k - a_{k-1} e^w] \\ &= [a_k e^w - a_{k+1} : a_{k-1} e^w - a_k], \end{aligned}$$

that is $\chi_n(\mathbb{H}_k) = G_n^{-k} \cdot \chi_n(\mathbb{H}_0)$. It is also clear that

$$\begin{cases} G_n^{-k} \cdot [1 : 0] = [-a_{-k-1} : -a_{-k}] = [a_{k+1} : a_k] \quad \text{and} \\ G_n^{-k} \cdot [0 : 1] = [a_{-k} : a_{-k+1}] = [a_k : a_{k-1}]. \end{cases}$$

Therefore χ_n wrap

- $\mathbb{H}_0$ on $\mathbb{C}\mathrm{P}^1$ around the axis from $[0 : 1]$ to $[1 : 0]$ and
- $\mathbb{H}_k$ on $\mathbb{C}\mathrm{P}^1$ around the axis from $[a_k : a_{k-1}]$ to $[a_{k+1} : a_k]$.

Note that (6) induces the following:

$$\lim_{k\to\infty} [a_k : a_{k-1}] = \lim_{k\to\infty} [a_{k+1} : a_k] = \left[1 : \frac{n - \sqrt{n^2-4}}{2}\right]$$

$$\lim_{k\to\infty} [a_{-k} : a_{-k-1}] = \lim_{k\to\infty} [a_{-k+1} : a_{-k}] = \left[1 : \frac{n + \sqrt{n^2-4}}{2}\right].$$

See Figure 5. We have thus proved that χ_n is a covering map if it is restricted onto

$$\mathcal{CP}_n = \mathbb{C}\mathrm{P}^1 \setminus \left\{ \begin{array}{l} [a_k : a_{k+1}] \ (k \in \mathbb{Z}), \\ \text{arc from } \left[1 : \frac{n-\sqrt{n^2-4}}{2}\right] \text{ to } \left[1 : \frac{n-\sqrt{n^2+4}}{2}\right] \end{array} \right\} \tag{10}$$

Now we have completed the proof of Theorem 1.1. □

Proof of Theorem 1.2. This is a corollary to Theorem 1.1. The inverse images of (8), (9), and (10) via $\pi : \mathbb{C}^2 \setminus \{(0,0)\} \to \mathbb{C}\mathrm{P}^1$ are given as follows:

$$\begin{aligned} &\pi^{-1}(\mathcal{CP}_1) = \mathbb{C}^2 \setminus \{z_1 = 0\} \cup \{z_2 = 0\} \cup \{z_1 = z_2\}, \\ &\pi^{-1}(\mathcal{CP}_2) = \mathbb{C}^2 \setminus \{z_2 = z_1\} \cup \{k z_2 = (k+1) z_1 \mid k \in \mathbb{Z}\}, \\ &\pi^{-1}(\mathcal{CP}_n) = \mathbb{C}^2 \setminus \{a_k z_2 = a_{k+1} z_1 \mid k \in \mathbb{Z}\} \\ &\qquad\qquad \cup \left\{ z_2 = \lambda z_1 \;\middle|\; \lambda \in \mathbb{R}, \frac{n - \sqrt{n^2-4}}{2} \le \lambda \le \frac{n + \sqrt{n^2-4}}{2} \right\}. \end{aligned}$$

Then the map $\mathcal{Z} : \mathrm{Stab}(P_n) \to \mathbb{C}^2$ is a covering map if it is restricted to each of these subspaces. □

References

[Bon89] A. I. Bondal. Representations of associative algebras and coherent sheaves. *Izv. Akad. Nauk SSSR Ser. Mat.*, 53(1):25–44, 1989.

[Bri07] Tom Bridgeland. Stability conditions on triangulated categories. *Ann. of Math. (2)*, 166(2):317–345, 2007.

[Bri09] Tom Bridgeland. Stability conditions and Kleinian singularities. *Int. Math. Res. Not. IMRN*, (21):4142–4157, 2009.

[CB92] William Crawley-Boevey. Exceptional sequences of representations of quivers. In *Proceedings of the Sixth International Conference on Representations of Algebras (Ottawa, ON, 1992)*, volume 14 of *Carleton-Ottawa Math. Lecture Note Ser.*, page 7, Ottawa, ON, 1992. Carleton Univ.

[IUU10] Akira Ishii, Kazushi Ueda, and Hokuto Uehara. Stability conditions on A_n-singularities. *J. Differential Geom.*, 84(1):87–126, 2010.

[Kra07] Henning Krause. Derived categories, resolutions, and brown representability. In *Interactions between homotopy theory and algebra*, volume 436 of *Contemp. Math.*, pages 101–139. Amer. Math. Soc., Providence, RI, 2007.

[Mac07] Emanuele Macrì. Stability conditions on curves. *Math. Res. Lett.*, 14(4):657–672, 2007.

[Oka07] So Okada. Stability conditions on projective spaces. *math.AG/0411220*, 2007.

[Rin94] Claus Michael Ringel. The braid group action on the set of exceptional sequences of a hereditary Artin algebra. In *Abelian group theory and related topics (Oberwolfach, 1993)*, volume 171 of *Contemp. Math.*, pages 339–352. Amer. Math. Soc., Providence, RI, 1994.

Department of Mathematical Sciences, Shinshu University, Matsumoto, Japan
E-mail address: `shiina@math.shinshu-u.ac.jp`